

## A Revision of the Collection of Noctuidae (Lepidoptera) reported from Korea by O. Herz in 1904

Vladimir S. KONONENKO

Institute of Biology and Pedology, Far Eastern Branch of the Russian Academy of Sciences,  
RF-690022 Vladivostok, Russia

---

**Abstract** The collection of Noctuidae reported by Herz (1904) from Korean the peninsula is revised and its corrected list is given. The identity of some species is confirmed or corrected with comments related to recent Korean faunistic literatures. The types of the noctuids described by Herz (1905) are revised and the lectotypes for seven taxa described are designated. New synonymies for two taxa (*Coryta grisea* Leech, 1900 = *Polydesma striata* Herz, 1904, **syn. n.**, *Apopestes indica* Moore = *Apopestes spectrum* var. *coreana* Herz, 1904, **syn. n.**) are proposed. The male genitalia of types and authenctical specimens for the taxa described by Herz are illustrated.

---

**Key words** Lepidoptera, Noctuidae, revision of collection, nomenclature, Korean peninsula

### INTRODUCTION

A well-known Russian lepidopterist, the conservator of Russian Imperial Zoological Museum, Otto Fedrovich Herz (1852–1905) was one of the first researcher of the Lepidoptera fauna in the Korean peninsula. He collected Lepidoptera in Korea by himself during the summer season 1884 at a small villiage Pung-Tung near Kimwah (38.5° N, 128° E), Kangwon Province. Herz also examined some materials collected by M. Jankowsky in 1894 at a valley of Tumankang river in the northern part of Korea. As a results of his study for the Korean Lepidoptera, Herz published "The list of Noctuidae and Geometridae from Korea" (Herz, 1904). It was a complete and accurate check list at that time, with numerous additions to previous publications (Leech, 1889, 1900) and with descriptions of nine new species, a variety, and a new genus of the Noctuidae.

Herz's materials was a part of collection of Great Prinz Nikolai Mikhailovich Romanov - an uncle of the last Russian Imperior Nikolai Romanov, famoust lepidoptera amateur, sponsor of many entomological expeditions and projects, a publisher of famous Russian lepidopterological issue "Memoirs sur Lepidopterous". Collection of N. M. Romanov was donated by him to Zoological Museum of Imperior Academy of Sciences in St. - Petersburg in 1900. At present these materials including Herz collection from Korea are preserved in the Zoological Institute of Russian Academy of Sciences. Like other

members of Imperior's family N. M. Romanov was shutted after revolution by bolsheviks in 1917 in Petrograd.

Since Herz's publication, during the subsequent 1890 years, the nomenclature and taxonomic status of many taxa for the Korean noctuids have been changed considerably. The present paper is based on the examination of Herz's materials and type specimens described by him from Korea, and dividing in three parts:

**1) A revised list of Herz's material.** In this part, the names of taxa are cited as same order and nomenclature as used by Herz in the left column. In the right column, the names of each taxon are arranged by the correct identification under the current status and nomenclature where they have been changed. The following abbreviations are used in the parenthesis: misid. – misidentification; missp. – misspelling; mispl. – misplacing, for the species treated in another genus (or under another generic name) in current sense; syn. – synonymic name at present, hom. – homonym; emend. – unnecessary emendation. Some taxa required more complete explanation are marked by indexes.

**2) Comments.** Comments to Herz's list contains explanations for some critical species corresponding to names used by him. In most cases, comments are given in relation to using Herz's data in Korean literatures on Noctuidae.

**3) A revised catalogue of the types described by Herz.** The taxa described by Herz's are indicated in the left column of Herz's list with asterisk (\*). In the catalogue, the names of species-group taxa are arranged in the alphabetical order with the following standard data: name of taxon, originally accepted taxonomic category (in round bracket): (sp) – species, (var) – variety; original combination with author and year of reference to the original description; type locality (TL); type material (TM). The data of labels are quoted exactly, separated by a stroke (/). The synonyms and current status for each taxa are given under the paragraph "ID" (identity) with references to corresponding literary sources.

The specimens in Herz' collection have small, printed standard labels with black frame: "Korea", on upper side "Herz" or "Korea Herz 1884" or "Korea" and "Jankowsky 1894". In addition all specimens have the labels in Russian: "Колл. Велл. Николая Михайловича." which means "Collection of Great Prince Nicolai Mikhailovich [Romanov]". The specimens from type series are supplied by blue or pinky label with "Origin". Some type specimens illustrated in photo plates in Herz's publication have the pins with upper part cut. Under description of a new of taxa Herz gave the label "Origin" to all specimens in type series, but did not designate as "Type" for a single specimen. Therefore the lectotypes are selected for seven taxa. In three cases the taxa were described, based on a single specimen, which is the holotype according to International Code of Zoological Nomenclature (ICNZ). For illustrations of genitalia, both types and authentional specimens identified by comparing with types were used.

## A REVISED LIST OF HERZ'S MATERIAL

### Original names by Herz (1904)

*Diphtera niveola* Motsh. (mispl.)

*Diphtera virens* Butl. (misid., mispl.)

*Diphtera alpium* Osb. var. *murrhina* Graes.

### Corrections (Current name)

– *Belciades niveola* (Motsch., 1866)

– *Belciana staudingeri* (Leech, 1900)

– *Moma kolthoffi* (Bryk, 1948)

- (misid., mispl.)
- Aplectoides caliginea* Butl. (mispl.) – *Anacronicta caliginea* (Butl., 1881)
- Acronicta major* Brem. – *Acronicta major* Brem., 1861
- Acronicta strigosa* F. – *Acronicta strigosa* (Den. & Schiff., 1775)
- Acronicta albonigra* Herz \*(mispl.) – *Hampsonidia albonigra* (Herz, 1904)
- Acronicta jankowskii* Obth. (mispl.) – *Hampsonidia jankowskii* (Obth., 1880)
- Acronicta tridens* D.& S. var. *incretata* Butl. (syn.) – *Acronicta intermedia* (Warr., 1909 )
- Acronicta cuspis* Hb. var. *leucocuspis* Butl. – *Acronicta leucocuspis* (Butl., 1878)
- Acronicta raphaelis* Obth. (emend.) – *Acronicta raphael* (Obth., 1884)
- Acronicta digna* Butl. – *Acronicta digna* (Butl., 1881)
- Acronicta rumicis* L. – *Acronicta rumicis* (L., 1758)
- Acronicta carbonaria* Graes. – *Acronicta carbonaria* (Graes., [1898] 1888)
- Acronicta lutea* Brem. – *Acronicta lutea* (Brem. & Grey, 1852)
- Agrotis obscura* Brahm. (syn., mispl.) – *Spaelotis ravida* (Den. & Schiff., 1775)
- Agrotis fuscostigma* Brem. (mispl.) – *Xestia fuscostigma* (Brem., 1861)
- Agrotis exusta* Butl. (mispl.) – *Sineugraphe exusta* (Butl., 1878)
- Agrotis exusta*, ab. *nigromaculata* Graes. (syn., mispl.) – *Sineugraphe bipartita* (Graes., [1889] 1888)
- Agrotis baja* F. (misid., mispl.) – *Xestia tabida* (Butl., 1878)
- Agrotis kollari* Led. (mispl.) – *Xestia kollari* (Led., 1853)
- Agrotis cissigma* Men. (mispl.) – *Lycophotia cissigma* (Men., 1859)
- Agrotis stupenda* Butl. (mispl.) – *Xestia stupenda* (Butl., 1878)
- Agrotis triangularis* Moore (mispl.) – *Albocosta triangularis* (Moore, 1867)
- Agrotis islandica* Stgr var. *rossica* Stgr. (mispl.) – *Euxoa ochrogaster rossica* (Stgr., 1881)
- Agrotis nigricostata* Stgr. (syn., mispl.) – *Agrotis tokionis* Butl., 1881
- Agrotis ypsilon* Rott. (emend., miss. auth.) – *Agrotis ipsilon* (Hfn., 1766)
- Agrotis segetum* D.& S. – *Agrotis segetum* (Den. & Schiff, 1775.
- Agrotis sibirica* Bsdv. – *Euxoa sibirica* (Bsdv., 1832)
- Mamestra adjuncta* Stgr. (syn., mispl.) – *Polia bombycina grisea* (Butl., 1878)
- Mamestra brassicae* L. – *Mamestra brassicae* (L., 1758)
- Mamestra splendens* Hb. (mispl.) – *Lacanobia splendens* (Hb., [1808])
- Dianthoecia cucubali* Fluess. (misid., mispl.) – *Sideridis honei* (Yoshimoto, 1989)
- Miana askoldis* Obth. (mispl.) – *Leucapamea askoldis* (Obth., 1880)
- Miana segregata* Butl. (mispl.) – *Nipponyx segregata* (Butl., 1878 )
- Miana variegata* Obth.(syn., mispl.) – *Pyrrhivalva sordida* (Butl., 1881)
- Miana strigilis* Cl. var. *an ab.* (mispl.) – *"Oligia" leuconephra* (Hmps., 1808 )
- Luperina depravata* Butl. (mispl.) – *Spodoptera depravata* (Butl., 1879)
- Hadena subpulchra* Alph. (mispl.) – *Euromioia subpulchra* (Alph., 1897)
- Hadena confecta* Walk. (misid., misspl.) – *Olivinebula oberthueri* (Stgr., 1892)
- Hadena funerea* Hein. (syn., mispl.) – *Apamea aquila oriens* (Warr., 1911)

- Hadena muscosa* Butl. (syn.) – *Atrachea nitens* (Butl., 1878)  
*Hadena jankowskii* Obth. (mispl.) – *Atrachea jankowskii* (Obth., 1879)  
*Hadena arcta* Led. (part., mispl.) – *Eucarta arcta* (Led., 1853)  
*Hadena arcta* Led. (part., misid., mispl.) – *Eucarta fasciata* (Butl., 1878)  
*Hadena arctides* Stgr. (mispl.) – *Eucarta arctides* (Stgr., 1888)  
*Dipterygia scabriuscula* L. (misid.) – *Dipterygia caliginosa* (Walk., 1858)  
*Rhisogramma saga* Butl. (mispl.) – *Auchmis saga* (Butl., 1878)  
*Antha grata* Butl. – *Antha grata* (Butl., 1881)  
*Chloantha intermedia* Brem. (emend., mispl.) – *Actinotia intermediata* (Brem., 1861)  
*Callopietria purpureofasciata* Piller (syn.) – *Callopietria juvenina* (Stoll., 1782)  
*Callopietria miracula* Herz\* (mispl.) – *Callopietria miracula* Herz, 1904 (mispl.)  
*Callopietria argyrosticta* Butl. – *Callopietria argyrosticta* (Butl., 1881)  
*Trachea atriplicis* L. – *Trachea atriplicis* (L., 1758)  
*Trachea melanospila* Koll. var. *lucia* Butl. (hom.) – *Trachea punkikonis* Mats., 1928 (*lucia* Butl., nec Feld. & Rghr., 1874)  
  
*Trachea kosakka* Obth. (syn.) – *Trachea melanospila* Koll., [1844]  
*Euplexia lucipara* L. – *Euplexia lucipara* (L., 1758)  
*Mania muscivirens* Butl. – *Mormo muscivirens* Butl., 1878  
*Orthogonia sera* Feld. – *Orthogonia sera* Feld. & Feld., 1862  
*Naenia contaminata* Walk. – *Naenia contaminata* (Walk., 1865)  
*Helotropha leucostigma* Hb. var. *laevis* Butl. (mispl.) – *Celaena leucostigma* (Hb., [1808])  
*Virgo amoena* Stgr. (syn.) – *Virgo datanidia* (Butl., 1885)  
*Nonagria graminea* Graes. (mispl.) – *Xylomoia graminea* (Graes., [1889] 1888)  
*Sesamia inferens* Walk. – *Sesamia inferens* (Walk., 1856)  
*Leucania impuncta* Stgr. (hom. mispl.) – *Mythimna simplex* (Leech, 1900), repl. name.  
*Leucania pallens* var. *melania* Stgr. (misident, mispl.) – *Mythimna pallens* (L., 1758)  
  
*Leucania velutina* Ev. (mispl.) – *Mythimna velutina* (Ev., 1846)  
*Leucania radiata* Brem. (mispl.) – *Mythimna radiata* (Brem., 1861)  
*Leucania albicosta* Moore (hom. misid., mispl.) – *Mythimna separata* (Walk., 1865)  
*Leucania unipuncta* Haw. (misid. mispl.) – *Mythimna separata* (Walk., 1865)  
*Leucania turca* L. (mispl.) – *Mythimna turca* (L., 1758)  
*Leucania divergens* Butl. (mispl.) – *Mythimna divergens* Butl., 1887  
*Leucania biundulata* Motsch. (syn. uncert.) – *Mythimna grandis* Butl., 1878  
*Leucania rufipennis* Butl. (mispl.) – *Mythimna rufipennis* Butl., 1878  
*Caradrina albosignata* Obth. et ab. *caeca* Obth. (mispl.) – *Athetis albisignatha* (Obth., 1879)  
  
*Caradrina lenta* Tr. var. *lentina* Stgr. (syn.) – *Athetis furvula* (Hb., [1808])  
*Hydrilla gluteosa* Tr. (mispl.) – *Athetis gluteosa* (Tr., 1835)  
*Hydrilla maurella* Stgr. (mispl.) – *Stygiodrina maurella* (Stgr., 1888)

- Hydrilla illustrata* Stgr. (mispl.)  
*Hydrilla tristis* Brem. ab. *lugens* Stgr. (syn.)  
*Amphipyra livida* F. var. *corvina* Motsch.  
*Amphipyra jankowskii* Obth.  
*Amphipyra erebina* Butl.  
*Amphipyra pyramidea* L. var. *monolitha* Gn.  
*Eutelia geyeri* F. & R.  
*Anuga constricta* Gn. (misid.)  
*Erastria argentula* Hb. var. *amurula* Stgr. (syn., mispl.)  
*Erastria oblitterata* Rmbr. (mispl.)  
*Erastria pusilla* View. (syn.)  
*Erastria nemorum* Obth. (mispl.)  
*Erastria fasciana* L. var. *guenei* Fall. (syn.)  
*Erastria fasciana* L. var. *stygia* Butl. (mispl.)  
*Erastria fasciana* L. var. *distinguenda* Stgr. (mispl.)  
*Erastria fentoni* Butl. var. *versicolor* Obth. (mispl.)  
*Rivula sericealis* Scop.  
*Milichia gemella* Leech (mispl.)  
*Ilattia stellata* Butl. (mispl.)  
*Amyna octo* Gn.  
*Emmelia trabealis* Scop.  
*Tarache bicoloria* Leech (mispl.)  
*Scoliopteryx libatrix* L.  
*Calpe capucina* Esp. (syn.)  
*Calpe aureola* Graes. (syn.)  
*Plusiodonta casta* Butl.  
*Telesilla amethystina* Hb. (mispl.)  
*Telesilla virgo* Tr. (mispl.)  
*Plusia agnata* Stgr. (mispl.)  
*Plusia ornatissima* Walk. (misid., mispl.)  
*Thermesia ussuriensis* Brem. (mispl.)  
*Remigia annetta* Butl. (mispl.)  
*Remigia archesia* Cram. (syn.)  
*Pelamia electaria* Brem. (mispl.)  
*Polydesma striata* Herz\* (syn. n.)  
*Zethes musculus* Men. (mispl.)  
*Zethes subfalcata* Men. (mispl.)  
*Ophiusa obscura* Brem. (mispl.)  
*Ophiusa coreana* Leech (mispl.)
- *Condica illustrata* (Stgr., 1888)  
 – *Athetis lepigone* (Mshl., 1907)  
 – *Amphipyra livida corvina* Motsch., 1866  
 – *Amphipyra jankowskii* Obth., 1884  
 – *Amphipyra erebina* Butl., 1878  
 – ?*Amphipyra monolitha* Gn., 1852 (id. uncert)  
 – *Eutelia geyeri* Feld. & Rog., 1774  
 – *Anuga multiplicans* Walk., 1858  
 – *Deltote bankiana* (F., 1775)  
 – *Phyllophila oblitterata* (Rmbr., 1833)  
 – *Pseudeustrotia candidula* (D. & S.)  
 – *Deltote nemorum* (Obth., 1880)  
 – *Protodeltote pydarga* (Hfn., 1766)  
 – ?*Sugia stygia* (Butl., 1878)  
 – *Protodeltote distinguenda* (Stgr., 1888)  
 – *Erastroides fentoni* (Obth., 1881)  
 – *Rivula sericealis* (Scop., 1763)  
 – *Dysmilichia gemella* (Leech, 1889)  
 – *Amyna stellata* (Butl., 1878)  
 – *Amyna octo* Gn., 1852  
 – *Emmelia trabealis* (Scop., 1763)  
 – *Acontia bicolora* (Leech, 1889)  
 – *Scoliopteryx libatrix* (L., 1758)  
 – *Calyptra thalictri* (Bkh., 1790)  
 – *Calyptra lata* (Butl., 1881)  
 – *Plusiodonta casta* Butl., 1878  
 – *Eucarta amethystina* (Hb., [1803])  
 – *Eucarta virgo* (Tr., 1835)  
 – *Acanthoplusia agnata* (Stgr., 1892)  
 – *Antoculeora locuples* (Obth., 1880)  
 – *Blasticorhinus ussuriensis* (Brem., 1861)  
 – *Mocis annetta* (Butl., 1878)  
 – *Mocis undata* (F., 1775)  
 – *Melapia electaria* (Brem., 1864)  
 – *Cortyta grisea* Leech, 1900  
 – *Arytrura musculus* (Men., 1852)  
 – *Arytrura subfalcata* (Men., 1852)  
 – *Dysgonia obscura* (Brem. & Grey, 1853)  
 – *Dysgonia coreana* (Leech, 1889)

- Ophiusa arcuata* Moore var. *mandshuriana* Stgr. (mispl.) – *Dysgonia mandshuriana* (Stgr., 1892)  
*Ophiusa annalis* Gn. (misid., mispl.) – *Dysgonia arctotaenia* Gn., 1852  
*Ophiusa dulcis* Butl. (mispl.) – *Dysgonia dulcis* (Butl., 1878)  
*Ophiusa dotata* F. (mispl.) – *Artena dotata* (F., 1794)  
*Spirama rectifasciata* Men. (mispl.) – *Metopta rectifasciata* (Men., 1863)  
*Sypna picta* Butl. (mispl.) – *Sypnoides picta* (Butl., 1877)  
*Sypna rectifasciata* Graes. (syn., mispl.) – *Sypnoides hercules* (Butl., 1881)  
*Edesena hamada* Feld. – *Edesena hamada* (Feld. & Rog., 1874)  
*Pseudophila amata* Brem & Grey (mispl.) – *Chrysorithrum amatum* (Brem. & Grey, 1853)  
*Pseudophila flavomaculata* Brem. (mispl.) – *Chrysorithrum flavomaculatum* (Brem., 1861)  
*Arcte coerulea* Gn. (mispl.) – *Arcte coerula* (Gn., 1852)  
*Lagoptera junio* Dalm. (mispl.) – *Thyas junio* (Dalm., 1823)  
*Ophideres tyrannus* Gn. (mispl.) – *Adris tyrannus* (Gn., 1852)  
*Catocala electa* Bkh. – *Catocala electa* Viev., 1790  
*Catocala nupta* L. var. *obscurata* Obth. – *Catocala nupta* (L., 1761)  
*Catocala fulminea* Scop. – *Catocala fulminea* (Scop., 1763)  
*Catocala koreana* Stgr. – *Catocala koreana* Stgr., 1892  
*Catocala abamita* Brem. & Grey – *Catocala abamita* Brem. & Grey, 1853  
*Catocala agitatrix* Graes. – *Catocala agitatrix* Graes., [1889] 1888  
*Catocala obscena* Alph. – *Catocala obscena* Alph., 1879  
*Catocala helena* Ev. – *Catocala helena* Ev., 1856  
*Catocala doerriesi* Stgr. – *Catocala doerriesi* Stgr., 1888  
*Catocala oblitterata* Men. (syn.) – *Catocala praegnax* Walk., [1858] 1859  
*Catocala puella* Leech. – *Catocala puella* Leech., 1889  
*Koraia pirata* Herz\* – *Catocala pirata* (Herz, 1904)  
*Apopestes spectrum* L. var. *koreana* Herz\* (syn.) – *Apopestes indica* (Moore, 1882)  
*Apopestes cataphanes* Hb. – *Autophila inconspicua* Butl., 1881  
var. *inconspicua* Butl. (mispl.)  
*Eccrita maxima* Brem. (mispl.) – *Lygephila maxima* (Brem., 1861)  
*Toxocampa recta* Brem. (mispl.) – *Lygephila recta* (Brem., 1864)  
*Toxocampa moellendorffi* Herz\* (mispl.) – *Lygephila moellendorffi* (Herz, 1904)  
*Aventiola pusilla* Butl. – *Aventiola pusilla* (Butl., 1879)  
*Egnasia pulcherima* Butl. (mispl.) – *Lophoruza pulcherrima* (Butl., 1879)  
*Epizeuxis lunulata* Herz\* (syn.) – *Idia curvipalpis* (Butl., 1879)  
*Capnodes lignea* Butl. (mispl.) – *Anatatha lignea* (Butl., 1879)  
*Capnodes cremata* Butl. (mispl.) – *Diomea cremata* (Butl., 1878)  
*Capnodes cinerea* Butl. (mispl.) – *Hypostrotia cinerea* (Butl., 1878)  
*Bleptina spacoalis* Walk. (mispl.) – *Bertula spacoalis* (Walk., 1859)  
*Simplicia rectalis* Ev. – *Simplicia rectalis* (Ev., 1842)

<i>Gisira signata</i> Butl. (mispl.)	– <i>Cidaripura signata</i> (Butl., 1879)
<i>Zanclognatha tarsipumalis</i> Hb.	– <i>Zanclognatha lunalis</i> (Scop., 1763)
<i>Zanclognatha tarsipennalis</i> Tr.	– <i>Zanclognatha tarsipennalis</i> (Tr., 1835)
<i>Zanclognatha tarsicrinalis</i> Knoch.	– <i>Herminia tarsicrinalis</i> (Knoch., 1782)
<i>Zanclognatha fumosa</i> Butl.	– <i>Zanclognatha fumosa</i> Butl., 1879
<i>Zanclognatha stramentacealis</i> Brem. (mispl.)	– <i>Herminia stramentacealis</i> Brem., 1864
<i>Zanclognatha umbrosalis</i> Stgr.	– <i>Zanclognatha umbrosalis</i> Stgr., 1892
<i>Zanclognatha trilinealis</i> Brem.	– <i>Paracolax trilinealis</i> (Brem., 1864)
<i>Zanclognatha bistrigata</i> Stgr. (mispl.)	– <i>Bertula bistrigata</i> (Stgr., 1888)
<i>Zanclognatha punctalis</i> Herz* (mispl.)	– <i>Sinarella punctalis</i> (Herz, 1904)
<i>Pangrapta vasava</i> Butl.	– <i>Pangrapta vasava</i> Butl., 1881
<i>Pangrapta flavomacula</i> Stgr.	– <i>Pangrapta flavomacula</i> Stgr., 1888
<i>Pangrapta suaveola</i> Stgr.	– <i>Pangrapta suaveola</i> Stgr., 1888
<i>Pangrapta indentalis</i> Leech	– <i>Pangrapta indentalis</i> Leech, 1888
<i>Egnasia costaemacula</i> Stgr. (mispl.)	– <i>Pangrapta costaemacula</i> Stgr., 1888
<i>Marmorinia obscurata</i> Butl. (mispl.)	– <i>Pangrapta obscurata</i> (Butl., 1888)
<i>Hydrillodes morosa</i> Butl.	– <i>Hydrillodes morosa</i> Butl., 1879
<i>Dierna timandra</i> Alph. (mispl.)	– <i>Naganoella timandra</i> (Alph., 1897)
<i>Gynaephila maculifera</i> Stgr.	– <i>Gynaephila maculifera</i> Stgr., 1892
<i>Madopa salicalis</i> D. & S. (mispl.)	– <i>Colobochyla salicalis</i> (Den. & Schiff., 1775)
<i>Madopa flavomacula</i> Obth. (mispl.)	– <i>Paragabara flavomacula</i> (Obth., 1888)
<i>Herminia gryphalis</i> H.–S. (mispl.)	– <i>Polypogon gryphalis</i> (H.–S., 1851)
<i>Herminia derivalis</i> Hb. (syn., mispl.)	– <i>Paracolax tristalis</i> (F., 1794)
<i>Herminia tentacularia</i> Hb. (mispl.)	– <i>Polypogon tentacularia</i> (Hb., 1758)
<i>Bomolocha nigrobasalis</i> Herz* (mispl.)	– <i>Hypena nigrobasalis</i> (Herz, 1904)
<i>Bomolocha stygiana</i> Butl. (mispl.)	– <i>Hypena stygiana</i> (Butl., 1878)
<i>Bomolocha bicoloralis</i> Graes. (mispl.)	– <i>Hypena bicoloralis</i> (Graes., [1889] 1888)
<i>Dichromia amica</i> Butl. (mispl.)	– <i>Hypena amica</i> (Butl., 1878)
<i>Hypena proboscidalis</i> L. var. <i>deleta</i> Stgr.	– <i>Hypena proboscidalis</i> (L., 1758)
<i>Hypena tristalis</i> Led.	– <i>Hypena tristalis</i> Led., 1853
<i>Hypenodes separatalis</i> Herz* (mispl.)	– <i>Schrankia separatalis</i> (Herz, 1904)

## COMMENTS

*Diphtera alpium* var. *murrhina*. – The species was misidentified by Herz as *Diphtera alpium* var. *murrhina* (Graes.), but it was described later as *Moma koltzoffi* (Bryk, 1948) (= *fulvicollis* de Lattin, 1948). For the cited synonymy, see Kononenko, 1996. The name *Moma orion* var. *murrhina* Graeser, [1889] 1888 is a junior synonym of *Moma alpium* (Osbeck, 1778).

*Agrotis baja* – At present, *Xestia baja* ([Denis & Schiffermüller], 1775) and *Xestia tabida* are treated

as two distinct species, forming a sister group since Boursin (1963). The former is distributed in the Western Palaearctic region, and the Far Eastern population belongs to the latter. However this specific pair needs a revision for clarifying their status as two distinct species or a subspecies. Author identified Korean population as *X. tabida*.

***Agrotis islandica* var. *rossica*** – In the current treatment, *Euxoa islandica* (Staudinger, 1857) is considered as a synonym of the Holarctic species *Euxoa ochrogaster* (Guenée, 1953) (Lafontaine, 1987). In the Western Palaearctic region, the latter represented as a subspecies *Euxoa ochrogaster* ssp. *islandica* (Staudinger, 1857) and in the Eastern Palaearctic region as *Euxoa ochrogaster* ssp. *rossica* (Staudinger, 1881).

***Dianthoecia cucubali*** – The species reported as *Dianthoecia cucubali* (a junior subjective synonym of *Sideridis (Aneda) rivularis* (Fabricius, 1775)) by Herz was reidentified as *Sideridis (Aneda) honeyi* Yoshimoto, 1989. Since Herz, this species had been reported in Korean publications as *Hadena rivularis*, *H. rivularis mandarina* (Leech, 1900), *H. cucubali* ([Denis & Schiffermüller], 1775) or *H. cucubali mandarina*. It was listed to the Check List of Insects from Korea (1994) as *Hadena rivularis*. Yoshimoto (1989) separated *Hadena mandarina* (Leech, 1900) from *rivularis* as a distinct species and described subspecies *H. rivularis honeyi*. The taxonomic status of the last was raised by Hacker (1996) to a good species. *S. rivularis*, *S. honeyi* and *S. mandarina* represent as a close sister species-group, which are hardly identifiable by the external appearance. Only one species, *S. honeyi* has been found in Korea, although all three species are distributed in the south of Russian Far East and N China (Hacker, 1996); the occurrence of other species in Korea is not impossible, but need a further investigation.

***Miana strigilis* var. an ab.** – The taxon reported as *Miana strigilis* Cl. var. an ab. [*Oligia strigilis* (Linnaeus, 1758)] by Herz was re-identified as "*Oligia leuconephra*. *Oligia strigilis* is a Western Palaearctic species and not distributed in the Far East. Since Herz, it was not reported for Korea or neighbouring countries. The generic position of "*Oligia leuconephra* is still unclear.

***Hadena subpulchra*** – The species was described from Korea by Alpheraky (1897) based on specimen collected by Jankowski from N Korea and reported for Korea by Leech (1900). Herz (1904) reported it as *Hadena subpulchra*. After these records it was referred under the genus *Hadena* in Korean literatures. Kononenko (1976) transferred *Hadena subpulchra* to the genus *Euromioia* (Subfam. Acronictinae) and reidentified *Euromioia subpulchra* as *E. mixta* Staudinger, 1892 by Kozhanchikov (1950). The species was also incorrectly listed in the Check List of Insects from Korea (1994) under combinations *Hadena subpulchra* and also *Euromioia subpulchra*.

***Hadena confecta*** – The species, reported by Herz [ZI, St. Petersburg] as *Hadena confecta*, was reidentified as *Olivinebula oberthueri* (Staudinger, 1892). This species was also misidentified by Leech (1900) and reported for Korea as *Triphaenopsis pulcherrima* Moore. Both species, *Olivinebula pulcherrima* (Moore, 1867) and *Olivinebula confecta* (Walker, 1858) are distinct from *O. oberthueri* and



are not represented in the Far East. Both names, *pulcherrima* and *confecta* were incorrectly used in Korean literatures. The name *Hadena confecta* was incorrectly spelled by Pak (1959) as *conferta*. This name as well as *pulcherrima* was listed in the Check List of Insects from Korea (1994) as "*Hadena conferta* [sic, misspelling] Walker" and *Triphaenopsis pulcherima* [sic] Moore. All three names (*pulcherrima*, *confecta* and *conferta*) related to *Olivinebula oberthueri* and should be excluded from the faunal list of the Korean Noctuidae.

*Hadena jankowskii* – The species was reported as *Hadena jankowskii* by Herz (1904). Since Herz, it has been incorrectly referred in Korean literatures as *Hadena jankowskii* or as *Trachea jankowskii* Obth. Under the last combination as well as *Atrachea jankowskii* Oberthür it was listed in the Check List of Insects from Korea (1994). The species was also incorrectly illustrated in Nam *et al.* (1982: Pl. 42, fig. 295a) as *Atrachea nitens* (Butler, 1878).

*Hadena arcta* – The material reported as *Hadena arcta* by Herz was reidentified as *Eucarta arcta* (Lederer, 1853) and partially as *Eucarta fasciata* (Butler, 1878).

*Dipterygia scabriuscula* – Herz misidentified Korean specimen of *Dipterygia caliginosa* as the Western Palearctic species, *Dipterygia scabriuscula*. Since Herz, this name has been cited in some Korean publications. The former species is not distributed in the Far East.

*Trachea melanospila* – The species reported as *Trachea melanospila* var. *lucia* by Herz is currently considered as *Trachea punkikonis* Matsumura, 1927, which is a replaced name for the secondary homonym, *Hadena lucia* Butler, 1878. Since Herz, it has been incorrectly referred in the Korean literature as *Trachea auriplena* (Walker, 1857) or as *T. lucilla* [sic] Butler [*T. lucia* Butler, 1878]. It was incorrectly listed in the Check List of Insects from Korea (1994) as *Trachea auriplena*, although this Indian-Himalayan species is not distributed in the Far East.

*Leucania albicosta* and *Leucania unipuncta* – The species reported as *Leucania unipuncta* and *Leucania albicosta* by Herz are the misidentification of *Mythimna (Pseudaletia) separata* Walker, 1865. The former species do not distributed in the Far East. *Mythimna (Pseudaletia) pallidicosta* (Hampson, 1894) is a replacement name for *Aletia albicosta* Moore, 1882 (homonym) applied to a distinct species. However, under the examination of Herzs material from Korea [ZIN, Spb.], this species was not found. Some specimens of *M. separata* (reported by Herz as *Leucania unipuncta*) have some reddish tint in the coloration of forewings, which make them somewhat similar to *M. pallidicosta*, but it lies in the limits of specific variability of *M. separata*. The name *albicosta* probably after Herz (1904) was repeated in Korean literature (Pak, 1959; Nam *et al.*, 1982) and listed in the Check List of Insects from Korea (1994) under a homonym name of *Pseudaletia albicosta*. The species *M. pallidicosta* (*albicosta*, hom.) is distributed in South East Asia, and migrating moths reaching to South of Japan. It was not found in Korea. Herzs record is probably due to the misidentification.

***Amphipyra pyramidea* var. *monolitha*** – The identity of Korean and continental Far Eastern population of *Amphipyra pyramidea/ monolitha* with N Indian species *Amphipyra monolitha* Guenee, 1852 is still unclear. A further examination for this taxon is needed.

***Anuga constricta***. – The species reported for Korea as *Anuga constricta* Guenee, 1852 by Herz is a misidentification of *Anuga multiplicans* (Walker). Herz incorrectly placed the name *multiplicans* in the synonymy of *constricta*. However these two taxa currently are treated as distinct species (Holloway, 1985; Poole, 1989). *A. constricta* is distributed in SE Asia, north to Vietnam and Thailand, and *A. multiplicans* has more wide distribution and reaches to China, Japan and Korea. Under this name the species was reported for Korea by Pak (1964: 54) and also by Park and Lee (1977: 5). Both names, *Anuga multiplicans* and *Anuga constricta* were listed in the Check List of Insects from Korea (1994). However, the name *Anuga constricta* should be excluded from Korean faunal list.


***Erastria fasciana* var. *stygia*** – The material reported by Herz was examined and reidentified as *P. pygarga*. The name *stygia* is currently applied to a distinct species *Sugia stygia* (Butler, 1878). It hardly distinguishable from very polymorphic *P. pygarga* by the external appearance,. The species *Sugia stygia* was reported in Korean literatures (Maruta 1929a; 1929b), probably on the base of earlier reports by Leech (1889; 1900) and Herz (1904). The occurrence of *Sugia stygia* is possible in Korea, but needs a further investigation.

***Plusia ornatissima*** – The records of *Antoculeora ornatissima* (Walker, 1858) by Herz and other authors from Korea and the Far East are due to misidentification of *Antoculeora locuples* (Oberthr, 1881). These taxa currently are considered as two distinct species (Kitching, 1987, Kononenko, 1990; Ronkay, 1996). *A. ornatissima* is a Indian-Himalayan species, and not distributed in the Far East. All records of *A. ornatissima* from the Far East may belong to *A. locuples*.

***Ophiusa annalis*** – The species reported for Korea (Herz, 1904: 43) as *Ophiusa analis* Gn. [*Dysgonia analis* (Guenee, 1852)] by Herz is a misidentification of *Dysgonia stuposa* (Fabricius, 1794). Material reported by Herz was examined and reidentified as *D. stuposa*. The name *D. analis* is incorrectly listed in the Check List of Insects from Korea (1994).

***Egnasia costaemacula*** – The identity of this species from Herzs collection is uncertain. The species *Pangrapta costaemacula* (Staudinger, 1888) forms a sister species-group with its allies recently described as *P. duplex* (Sugi & Kononenko, 1996). The latter is the most common species of *Pangrapta* in Korea, Japan and China. During the examination of Herz's material, the specimen identified as *costaemacula* was not dissected, however, the occurrence of *costaemacula* in Korea, is quite possible in its northern part at least.

***Sypna rectifasciata*** – Material reported as *Sypna rectifasciata* by Herz was reidentified as *Sypnoides hercules* (Butler, 1881). The name *rectifasciata* is a junior synonym of *hercules*. Both name



**Figs 1-4.** Male genitalia: 1. *Hampsonia albonigra* (Herz), Russian Far East, Primorye terr., authentic specimen; 2. *Apopetes indica* (Moore) (= *Apopetes spectrum* var. *koreana* Herz), Russian Far East, Primorye, authentic specimen; 3. "*Callopietria*" *miracula* Herz, Korea, holotype; 4. *Lygephila moellendorffi* (Herz), Korea, paralectotype.

is listed in the Check List of Insects from Korea (1994) as *Sypna rectifasciata* and *Sypnoides hercules*. The former should be excluded from the faunal list of Korean Noctuidae as a synonym.

***Catocala puella*** Leech – Author could not find specimens in the collection of ZI, St. Petersburg, which was collected from Korea and identified by Herz. The species *Catocala puella* Leech, 1889, described from Korea, is known only by the type, preserved in BMNH, London, and also by two documented records in N and S Korea.

#### A REVISED CATALOGUE OF THE TYPES

***albonigra*** (sp) – *Acronicta albonigra* Herz, 1904: 269, pl.1, fig. 3. TL: Korea, Pun-tung villiage

[38.5° N, 128° E]. TM: Lectotype (hereby designated): male with labels: "Korea/Herz/Orig./**Колл. Вел. Николая Михайловича**". Genit. prep. N 17749. Paralectotypes: 6 males with the same labels, one male with label "Prep. 3487".

ID: *Hampsonia albonigra* (Herz, 1904), a good species (Subf. Acronictinae).

Remarks. A distinct species, distributed in Korea, Japan, Russian Far East. In current treatment it is placed to the genus *Craniophora* (Sugi, 1892). However, author treats *Hampsonia* (type species *Apatele jankowskii* Oberthür, 1880) and *Craniophora* (type species *Noctua ligustri* [Denis & Schiffermüller], 1775) as a distinct genera. Male genitalia for the authentical specimen of *Hampsonia albonigra* (Herz, 1904) is illustrated in the fig. 1.

**koreana (var)** – *Apopestes spectrum* L. var. *koreana* Herz, 1904: 315, pl.1, fig. 11. TL: North Korea [Tumangang valley]. TM: Lectotype (hereby designated): female with label: "Jankowsky, 1894 /Korea/ Orig./**Колл. Вел. Николая Михайловича**". The pin of the lectotype is cut at upper part. The specimen was illustrated by Herz. Paralectotypes: 5 ♂, 5 ♀, with same labels; 1 ♀: "Korea / Herz / Orig./**Колл. Вел. Николая**" and red label "Sp. *Spectrum* L. var. *phantasma* Ev."

ID: *Apopestes indica* (Moore, 1883) (= *Apopestes spectrum* L. var. *koreana* Herz, **syn. n.**) (Subf. Ophiderinae).


Remarks. The species was described by Herz from North Korea as *Apopestes spectrum* var. *koreana* Herz. Its status was revised by Boursin (1955) to a good species. Later Boursin put the name *koreana* to a synonymy of *Apopestes indica* Moore, 1883 in his unpublished Check List of Palaearctic Noctuidae, but it was not officially synonymized. The types of *Apopestes indica* are conspecific with *koreana* (L. Ronkay, pers. comm.). Poole (1989) incorrectly referred *Apopestes indica* as a synonym of *Apopestes spectrum* (Esper, 1787) and *koreana* as a good species. Both names *Apopestes koreana* and *Apopestes spectrum* were incorrectly included in the Check List of Insects from Korea (1994). The species *A. spectrum* does not distributed in the Far East, the name *koreana* is a synonym of *Apopestes indica*. The genitalia of the authentical specimen, compared with types *koreana*, is illustrated in the fig. 2.

**miracula (sp)** – *Calloplistria miracula* Herz, 1904: 284, pl.1, fig. 10. TL: Korea, Pun-tung villiage [38.5° N, 128° E]. TM: Holotype: a male with labels: "Korea / Herz /Orig./**Колл. Вел. Николая Михайловича**."

ID: "*Calloplistria*" **miracula** Herz, 1904, a good species (Subf. Amphipyryinae).

Remarks. A distinct specimen was found in the recent Korean collection. Its generic position is uncertain. Herz placed it to *Calloplistria*, however, its external morphology and male genitalia do not correspond to *Calloplistria juvenina* (Stoll, 1782), the type species of the genus. Apparently this species belongs to lower Amphipyryinae. Until now the species is known only from Korea. Male genitalia of the holotype of *Calloplistria miracula* Herz, 1904, is illustrated in the fig. 3.

**moellendorffi (sp)** – *Toxocampa moellendorffi* Herz, 1904: 318, pl.1, fig. 1. TL: Korea, Pun-tung villiage [38.5° N, 128° E]. TM: Lectotype (hereby designated): male, "Korea / Herz / Orig./**Колл. Вел. Николая Михайловича**". The specimen was illustrated by Herz, the upper part of pin is cut.



**Figs 5-8.** Male genitalia: 5. *Idia curvipalpis* (Butler) (= *Epizeuxis lunulata* Herz), Russian Far East, Primorye, authentic specimen; 6. *Hypena nigrobasalis* (Herz), Korea, holotype; 7. *Catocala pirata* (Herz), Russian Far East, Primorye, authentic specimen; 8. *Sinarella punctalis* (Herz), Russian Far East, Primorye, authentic specimen.


Paralectotype: 1 ♂ with same labels.

ID: *Lygephila moellendorffi* (Herz, 1904), a good species (Subf. Ophiderinae).

Remarks. A good species, however, No additional record after Herz's description is appeared. Male genitalia of the paralectotype is illustrated in the fig. 4.

*lunulata* (sp) – *Epizeuxis* (*Helia*) *lunulata* Herz, 1904: 320, pl.1, fig. 12. TL: Korea, Pun-tung villiage [38.5° N, 128° E]. TM: Lectotype (hereby designated): a male, "Korea / Herz / Orig. [red label]/ Колл. Вел. Николая Михайловича". Paralectotypes: 1 ♂ (without abdomen) with same label.

ID: *Idia curvipalpis* (Butler 1879) (Subf. Herminiinae).



**Figs 9-12.** Male genitalia: 9. *Schrankia separatalis* (Herz), Russian Far East, Primorye, authentic specimen; 10. *Cortyta grisea* (Leech) (= *Polidesma striata* Herz), Korea, authentic specimen. adults of types; 11. "*Callopietria*" *miracula* Herz, Korea, holotype 12. *Lygephila moellendorffi* (Herz), Korea, paralectotype.

Remarks. The name is currently considered to be a junior synonym of *Idia curvipalpis* (Butler), synonymized by Remm & Martin (1979). The species was illustrated by Owada (1982). Male genitalia of the authentic specimen is illustrated in the fig. 5.

**nigrobasalis (sp)** – *Bomolocha nigrobasalis* Herz, 1904: 329, pl.1, fig. 13. TL: Korea, Pun-tung villiage [38.5° N, 128° E]. TM: Lectotype (hereby designated): a male, "Korea Herz 1884 / Orig./ Колл. Вел. Николая Михайловича," genitalia preparete No 17490. Paralectotypes: 2 ♂, with labels: "Jankowsky, 1894 /Korea/ Orig./ Колл. Вел. Николая Михайловича".

ID: *Hypena nigrobasalis* (Herz, 1904), a good species (Subf. Hypeninae).

Remarks. Currently considered as a good species. The species was illustrated by Sugi (1982). Male genitalia of the lectotype is illustrated in the fig. 6.

**pirata (sp)** – *Koraia pirata* Herz, 1904: 314, pl.1, fig. 18. TL: Korea, Pun-tung villiage [38.5° N,

128° E]. TM: Lectotype (hereby designated): 1 ♀ with labels: "Korea / Herz / Orig./**Колл. Вел. Николая Михайловича**". The pin of the lectotype is cut at upper part. The specimen was illustrated by Herz. Paralectotype: a female with the same labels and additional label "Microprepare N 14652".

ID: *Catocala pirata* (Herz, 1904), a good species (Subf. Catocalinae).

Remarks. The genus *Koraia* Herz described by Herz (1904) on the same publication was synonymised with *Catocala* Schrank by Kononenko (1990: 29, note 86). Male genitalia of the authentical specimen to compare with types is illustrated in the fig. 7.

*punctalis* (sp) – *Zanclognatha punctalis* Herz, 1904: 324, pl.1, fig. 9. TL: Korea, Pun-tung villiage [38.5° N, 128° E]. TM: Holotype: a female with labels: "Korea / Herz / Orig./**Колл. Вел. Николая Михайловича**".

ID: *Sinarella punctalis* (Herz, 1904) (Subf. Herminiinae).

Remarks. Currently considered as a good species. It was illustrated by Owada (1982; 1987). Male genitalia was illustrated by Yamamoto & Sugi (1955) as *Zanclognatha nigrobasalis* Yamamoto & Sugi (junior synonym of *punctalis*), Remm & Martin (1987) and also by Owada. Male genitalia of the authentical specimen is illustrated in the fig. 8.

*separatalis* (sp) – *Hypenodes separatalis* Herz, 1904: 331, pl.1, fig. 7. TL: Korea, Pun-tung williage [38.5° N, 128° E]. TM: Lectotype (hereby designated): a male with labels "Korea / Herz / Orig./**Колл. Вел. Николая Михайловича**". Paralectotypes: 2 ♂, 1 ♀, with the same labels.

ID: *Schrankia separatalis* (Herz, 1904) (Subf. Hypenodinae).

Remarks. Currently considered as a good species. It was illustrated by Sugi (1982). Male genitalia (fig. 9) was illustrated by Remm & Martin (1979).

*striata* (sp) – *Polydesma striata* Herz, 1904: 303, pl.1, fig. 4. TL: North Korea [?Tumanghan valley]. TM: Holotype: male with labels: "Jankowsky 1894/ Orig./**Колл. Вел. Николая Михайловича**".

ID: *Polydesma striata* Herz, 1904 is a junior synonym (syn. n.) of *Cortyta grisea* (Leech, 1900) (Subf. Catocalinae).

Remarks. The type specimen is identical with *Cortyta grisea* (Leech, 1900, described from China (Trans. ent. Soc. Lond. 1900: 552, *Polydesma*). The latter taxon was illustrated by Hampson (1913, text fig. 82) and Chen (1982). The new synonymy was established, based on the comparision of imagos and male genitalia of the holotype of *Polydesma striata* Herz with specimen from Korea which was identified as *Cortyta grisea* (Leech, 1900). In the original description, Herz incorrectly indicated the type as a female but the holotype of *P. striata* is a male. The specimen is in some worn condition, therefore wing pattern is not contrasted as fresh ones. Male genitalia of the holotype of *Polygesma striata* Herz is illustrated in the fig. 10.

## ACKNOWLEDGMENTS

I wish to express my cordial thanks to Dr. A. L. Lvovsky and Dr. I. L. Sukhareva, Zoological Institute, Russian Academy of Sciences, St.-Peterburg for their kind consultation and help during my study, with supplying the collection under their curation and preparation of the present article. I am deeply grateful to Dr. S. B. Ahn (Agricultural Sciences and Technology Institute, Rural Development Administration, Suwon, Korea) and Prof. K. T. Park (Center for Insect Systematics, Chuncheon, Korea) for their critical review of the manuscript.

## REFERENCES

- Alpheraky, S. 1897. Lépidoptères our et de la Am Corée. In M. Romanoff, *Mem. Lépid.* 9: 151–184.
- Boursin, Ch. 1955. Die *Apopestes* Hb.-und *Autophila* Hb.-Arten aus Dr. h. c. Höne's China-Ausbeuten. *Z. Wien. ent. Ges.* 40: 164–171.
- Boursin, Ch. 1963. Die Noctuidae-Arten aus Dr. h. c. Hne's China-Ausbeuten. V. Der Amathes-Komplex; VI Die Gattungen: *Raddea* Alph., *Estimata* Kozh., *Erebophasma* n. gen., *Spinipalpa* Alph. und *Perissandra* Warren; VII. Die Gattung *Ochropleura* Hb. *Forsch. Ber. Landes Nordrhein-Westfalen.* 1170: 1–107, pls 1–22, 1 map.
- Ent. Soc. Korea & Korean Soc. Appl. Ent. 1994. *Check List of Insects from Korea*. Kon-Kuk Univ. Press. Seoul. 744 pp.
- Chen, Y. 1982. Noctuidae. *Iconographia Insectorum Sinicorum*. Beijin. Science press. III: 237–390+1–24, pls 76–118.
- Nam, S.H. 1982. Noctuidae. In Kim *et al.*, *Illustrated Flora and Fauna of Korea*. Vol. 26. Insecta (8). 919 pp [in Korean with English synonymic list].
- Hacker, H. 1996. Revision der Gattung *Hadena* Schrank, 1802 (Lepidoptera: Noctuidae). *Esperiana* 5: 7–687.
- Hampson, G.F. 1913. *Catalogue of the Lepidoptera Phalaenae in the British Museum*. Vol. 12: i–xiii, 1–858 (1913); London, Taylor and Francis.
- Herz, O. 1904. Lepidoptera von Korea. Noctuidae & Geometridae. *Ann. Mus. Zool. Acad. imp. Sci. St.-Petersbourg* 9: 263–390, pl. 1.
- Holloway, J.D. 1985. The moths of Borneo Noctuidae: Euteliinae, Stictopterinae, Plusiinae, Pantheinae. *Malay. Nat. J.* 38 [Separately published as Moths of Borneo 14]: 157–317.
- Kitching, I.J. 1987. Spectacles and Silver Ys: a synthesis of the systematics, cladistic and biology of the Plusiinae (Lepidoptera: Noctuidae). *Bull. Br. Mus. nat. Hist. (Ent.)* 54: 75–261.
- Kononenko, V.S. 1976. Little known species of moths (Lepidoptera, Noctuidae) from the southern part of the Maritime Province. *Trudy zool. Inst., Leningr.* 67: 61–67. (In Russian).
- Kononenko, V.S. 1990. Synonymic Check List of the Noctuidae (Lepidoptera) of the Primorye Territory, the Far East of U. S. S. R. *Tinea* 13, Suppl. 40 pp.
- Kononenko, V.S. 1996. A revised catalogue of types of the Noctuidae (Lepidoptera) described by F. Bryk (1948) from the Korean Peninsula. *Ins. Koreana* 13: 1–26.
- Kozhanchikov, I. 1950. *Fauna SSSR. Nasekomye Cheshuekrylye* [Insecta, Lepidoptera]. Vol. 12. Volnyanki (Orgiidae). Leningrad. 581 pp.
- Lafontaine, J.D. 1987. Noctuoidea: Noctuidae (Part) Noctuinae (Part – Euxoa). In R. B. Dominik *et al.*, *Moths*



*of America North of Mexico* 27(2): 1–237.

Leech, J.H. 1889. On the Lepidoptera of Japan and Korea. Part III. Heterocera, sect. II. *Proc. zool. Soc. Lond.* 1889: 474–571.

Leech, J.H. 1900. Lepidoptera Heterocera from northern China, Japan and Corea. Part. 3. *Trans. ent. Soc. Lond.* 1900: 9–161; Part 4. *Ibidem.* 1900: 511–663.

Maruta, S.T. 1929a. A List of Lepidoptera in Kyojo and vicinity, *Bull. Agr. Exp. St. Govt., Chosen* 4(2): 131

Maruta, S.T. 1929b. Nocturnal flying insect. *Ann. Agr. Exp. St. Govt., Chosen* 4(6): 543.

Owada, M. 1982. Noctuidae (Herminiinae). In Inoue H., S. Sugi, H. Kuroko, S. Moriuti, and A. Kawabe. *Moths of Japan* 1: 913–935 ; 2: 405–408, pls 224–226, 356, 381–392. Kodansha, Tokyo.

Owada, M. 1987. A taxonomic study of the subfamily Herminiinae of Japan. *Natn. Sci. Mus. Tokyo*: 1–208. pp.

Pak, S.W. 1959. A Hand List of Korean Moths. *For. Exp. Sta. Korea*, 1: 25–26.

Park, K.T. and S.M. Lee. 1977. Description of one new species and notes on forty-three unrecorded species of Noctuidae (Lepidoptera) from Korea. *Korean J. Ent.* 7 (2): 1–10 [in Korean with English abstract].

Poole, R.W. 1989. Noctuidae. *Lepidopterorum Catalogues (New Series)*. Fascicle 118, 1314 pp. E. J. Brill, Leiden.

Remm, H.J. and M. Martin. 1979. On the morphology and systematic of the Hypeninae (Lepidoptera, Noctuidae). *Uchen. Zap. Tartu gos. Univ.* (483): 134–163.

Ronkay, L. 1996. A new species of *Antoculeora Ichinose* (Lepidoptera, Noctuidae, Plusiinae) from Taiwan. *Bull. Natn. Sci. Mus., Tokyo, Ser. A*, 23(1), p. 35–49.

Sugi, S. 1982. Noctuidae (except Herminiinae). In Inoue H., S. Sugi, H. Kuroko, S. Moriuti, and A. Kawabe. *Moths of Japan* 1: 669–913, 2: 334–405, pls. 37, 164–223, 229, 278, 355–280. Kodansha, Tokyo.

Yamamoto, Y. and S. Sugi. 1955. Early stage of two Herminiinae moths on moss, including a new species (Lepidoptera, Noctuidae). *Tinea* 2(1/2): 109–113.

Yoshimoto, H. 1989. *Hadena mandarina* and *H. rivularis* (Lepidoptera, Noctuidae), with description of a new subspecies and notes on a synonymy. *Japan Heterocerists' J.* (155): 73–76.

## Herz(1904)에 의해 한반도에서 보고된 밤나방과의 재정리

V.S. KONONENKO

러시아과학원극동분소 생물토양연구소, RF-690000 블라디보스톡, 러시아

1904년 Herz에 의해 한반도에서 보고된 밤나방과 種들을 재정리하고, 수정한 학명의 種목록을 보고한다. 또한 그에 의해 誤同定으로 발표되었던 種들에 대해 설명과 함께 재동정 결과를 정리하였으며, 일부 種의 모식표본 정리와 함께 種에 대한 Lectotype을 설정하였다. 두 중, *striata* Herz와 *coreana* Herz을 각각 *Coryta grisea* Leech와 *Apopestes indica* Moore의 synonym으로 정리하고, 몇몇 주요 種에 대해 성충의 사진과 수컷생식기를 도해하였다.

검색어 : 나비목, 밤나방과, 재정리, 명명규약, 한반도

(Received: November 5, 1997)

(Accepted: September 10, 1998)